



Farmer's Friend Horse Mineral & Vitamin Block

For Horses on Pasture

Guaranteed Analysis

| | |
|------------------------|---------------|
| Calcium (Min) | 12.50% |
| Calcium (Max) | 15.00% |
| Phosphorus (Min) | 7.00% |
| Salt (Min) | 13.90% |
| Salt (Max) | 16.60% |
| Cobalt (Min) | 1.1 ppm |
| Copper (Min)..... | 150 ppm |
| Iron (Min) | 2,240 ppm |
| Manganese (Min)..... | 250 ppm |
| Selenium (Min) | 21.1 ppm |
| Zinc (Min)..... | 500 ppm |
| Vitamin A (Min) | 100,000 IU/lb |
| Vitamin D-3 (Min)..... | 10,000 IU/lb |
| Vitamin E (Min) | 1,000 IU/lb |

CAUTION: Use as directed. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

Manufactured By:
Ridley Block Operations
Mankato, MN 56001

Ingredients

Monocalcium Phosphate, Dicalcium Phosphate, Processed Grain By-Products, Calcium Carbonate, Salt, Molasses Products, Calcium Hydroxide, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement and Mineral Oil.

INTRODUCTORY PERIOD

Position the blocks 50 to 75 feet from loafing, feeding and watering areas during the first week of feeding. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of the blocks does not correct the situation, remove the blocks and evaluate the overall feeding program.

NET WEIGHT: 40 LB (18.14 KG)

Feeding and Management

1. Feed at a rate of one block per 5 to 15 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to prevent crowding - 10 feet minimum.
3. Do not allow animals to run out of blocks. When a block is two-thirds consumed, provide a new block near it. Place very small pieces in a feed trough to be cleaned up.
4. Horses should consume an average of 2.5 to 5 ounces per head daily.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. Feed as the sole self-fed source of salt, minerals or vitamins.

For additional information, please contact your Triple Trust® representative.

Batch #

10346

TT1404
0216



Farmer's Friend Horse Mineral & Vitamin Block

For Horses on Pasture

Guaranteed Analysis

| | |
|------------------------|---------------|
| Calcium (Min) | 12.50% |
| Calcium (Max) | 15.00% |
| Phosphorus (Min) | 7.00% |
| Salt (Min) | 13.90% |
| Salt (Max) | 16.60% |
| Cobalt (Min) | 1.1 ppm |
| Copper (Min)..... | 150 ppm |
| Iron (Min) | 2,240 ppm |
| Manganese (Min)..... | 250 ppm |
| Selenium (Min) | 21.1 ppm |
| Zinc (Min)..... | 500 ppm |
| Vitamin A (Min) | 100,000 IU/lb |
| Vitamin D-3 (Min)..... | 10,000 IU/lb |
| Vitamin E (Min) | 1,000 IU/lb |

CAUTION: Use as directed. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

Manufactured By:
Ridley Block Operations
Mankato, MN 56001

Ingredients

Monocalcium Phosphate, Dicalcium Phosphate, Processed Grain By-Products, Calcium Carbonate, Salt, Molasses Products, Calcium Hydroxide, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement and Mineral Oil.

INTRODUCTORY PERIOD

Position the blocks 50 to 75 feet from loafing, feeding and watering areas during the first week of feeding. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of the blocks does not correct the situation, remove the blocks and evaluate the overall feeding program.

NET WEIGHT: 40 LB (18.14 KG)

Feeding and Management

1. Feed at a rate of one block per 5 to 15 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to prevent crowding - 10 feet minimum.
3. Do not allow animals to run out of blocks. When a block is two-thirds consumed, provide a new block near it. Place very small pieces in a feed trough to be cleaned up.
4. Horses should consume an average of 2.5 to 5 ounces per head daily.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. Feed as the sole self-fed source of salt, minerals or vitamins.

For additional information, please contact your Triple Trust® representative.

Batch #

10346

TT1404
0216