



14:4 Rumensin® 1200 Mineral

MEDICATED

For Further Manufacture of Feed

For increased rate of weight gain and for prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

Active Drug Ingredient

Monensin 1,200 g/ton

Guaranteed Analysis

Calcium (Min)	12.70%
Calcium (Max)	15.20%
Phosphorus (Min)	4.00%
Salt (Min)	18.20%
Salt (Max)	21.80%
Magnesium (Min)	2.00%
Copper (Min)	1,500 ppm
Manganese (Min)	4,000 ppm
Selenium (Min)	11 ppm
Zinc (Min)	4,000 ppm
Vitamin A (Min)	200,000 IU/lb
Vitamin D-3 (Min)	37,500 IU/lb
Vitamin E (Min)	100 IU/lb

Ingredients

Calcium Carbonate, Salt, Monocalcium Phosphate, Dicalcium Phosphate, Processed Grain By-Products, Magnesium Oxide, Molasses Products, Hydrogenated Soybean Oil, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylendiamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil and Red Iron Oxide.

Instruction for Mixing and Feeding

For increased rate of weight gain: Thoroughly mix 666 lbs. of Herdbuilder™ 14:4 Rumensin® 1200 Mineral with 1,334 lbs. of non-medicated feed. Feed 1 (one) pound of this medicated mixture to supply 199 mg of monensin. Cattle should receive not less than 50 mg nor more than 200 mg monensin per head per day in not less than 1 lb. of feed or, after the fifth day at the rate of 400 mg per head per day every other day in not less than 2 lb. of feed for the state drug claim. During the first 5 days of feeding, cattle should receive no more than 100 mg per day contained in not less than 1 lb. of feed. Thoroughly mix 333 lbs. of Herdbuilder™ 14:4 Rumensin® 1200 Mineral with 1,667 lbs. of non-medicated feed. Feed 1 (one) pound of this medicated mixture to supply 100 mg of monensin.

SEE RESERVE SIDE FOR ADDITIONAL FEEDING DIRECTIONS AND PRECAUTIONARY STATEMENTS.

NET WEIGHT: 50 lb (22.7 kg)

Batch #

10368

TT2303

1216

Rumensin® is a registered trademark of Elanco.
Herdbuilder™ is a trademark of Central Farm Supply of Kentucky.
Triple Trust® is a registered trademark of Central Farm Supply of Kentucky.
Manufactured By:
Ridley Block Operations, Mankato, MN 56001



14:4 Rumensin® 1200 Mineral

MEDICATED

For Further Manufacture of Feed

For increased rate of weight gain and for prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

Active Drug Ingredient

Monensin 1,200 g/ton

Guaranteed Analysis

Calcium (Min)	12.70%
Calcium (Max)	15.20%
Phosphorus (Min)	4.00%
Salt (Min)	18.20%
Salt (Max)	21.80%
Magnesium (Min)	2.00%
Copper (Min)	1,500 ppm
Manganese (Min)	4,000 ppm
Selenium (Min)	11 ppm
Zinc (Min)	4,000 ppm
Vitamin A (Min)	200,000 IU/lb
Vitamin D-3 (Min)	37,500 IU/lb
Vitamin E (Min)	100 IU/lb

Ingredients

Calcium Carbonate, Salt, Monocalcium Phosphate, Dicalcium Phosphate, Processed Grain By-Products, Magnesium Oxide, Molasses Products, Hydrogenated Soybean Oil, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylendiamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil and Red Iron Oxide.

Instruction for Mixing and Feeding

For increased rate of weight gain: Thoroughly mix 666 lbs. of Herdbuilder™ 14:4 Rumensin® 1200 Mineral with 1,334 lbs. of non-medicated feed. Feed 1 (one) pound of this medicated mixture to supply 199 mg of monensin. Cattle should receive not less than 50 mg nor more than 200 mg monensin per head per day in not less than 1 lb. of feed or, after the fifth day at the rate of 400 mg per head per day every other day in not less than 2 lb. of feed for the state drug claim. During the first 5 days of feeding, cattle should receive no more than 100 mg per day contained in not less than 1 lb. of feed. Thoroughly mix 333 lbs. of Herdbuilder™ 14:4 Rumensin® 1200 Mineral with 1,667 lbs. of non-medicated feed. Feed 1 (one) pound of this medicated mixture to supply 100 mg of monensin.

SEE RESERVE SIDE FOR ADDITIONAL FEEDING DIRECTIONS AND PRECAUTIONARY STATEMENTS.

NET WEIGHT: 50 lb (22.7 kg)

Batch #

10368

TT2303

1216

Rumensin® is a registered trademark of Elanco.
Herdbuilder™ is a trademark of Central Farm Supply of Kentucky.
Triple Trust® is a registered trademark of Central Farm Supply of Kentucky.
Manufactured By:
Ridley Block Operations, Mankato, MN 56001

Instruction for Mixing and Feeding

For coccidiosis control and prevention: Thoroughly mix Herdbuilder™ 14:4 Rumensin® 1200 Mineral with non-medicated feed as indicated in the table below. Cattle should receive 0.14 to 0.42 mg monensin per pound of bodyweight per day up to 200 mg/head/day depending on the severity of challenge. During the first 5 days of feeding, cattle should receive no more than 100 mg per day contained in not less than 1 lb. of feed.

Recommended Feeding Level	Desired Intake of Monensin	Herdbuilder™ 14:4 R1200 Mineral	Non-Medicated Feed
lbs/hd/d	mg/hd/d	lbs/ton	lbs/ton
2	50	83.25	1,916.75
2	60	100	1,900
2	100	166.5	1,833.5
2	200	333	1,667
1	150	500	1,500
1	200	666	1,334

For additional information contact your Herdbuilder™ representative.

CAUTION: Use as Directed. Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by equines has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

NET WEIGHT: 50 lb (22.7 kg)

Batch #

10368

TT2303

1216

Manufactured By:
Ridley Block Operations, Mankato, MN 56001

Instruction for Mixing and Feeding

For coccidiosis control and prevention: Thoroughly mix Herdbuilder™ 14:4 Rumensin® 1200 Mineral with non-medicated feed as indicated in the table below. Cattle should receive 0.14 to 0.42 mg monensin per pound of bodyweight per day up to 200 mg/head/day depending on the severity of challenge. During the first 5 days of feeding, cattle should receive no more than 100 mg per day contained in not less than 1 lb. of feed.

Recommended Feeding Level	Desired Intake of Monensin	Herdbuilder™ 14:4 R1200 Mineral	Non-Medicated Feed
lbs/hd/d	mg/hd/d	lbs/ton	lbs/ton
2	50	83.25	1,916.75
2	60	100	1,900
2	100	166.5	1,833.5
2	200	333	1,667
1	150	500	1,500
1	200	666	1,334

For additional information contact your Herdbuilder™ representative.

CAUTION: Use as Directed. Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by equines has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

NET WEIGHT: 50 lb (22.7 kg)

Batch #

10368

TT2303

1216

Manufactured By:
Ridley Block Operations, Mankato, MN 56001