

UltraPro[™] 20% AN Block

GUARANTEED ANALYSIS

| Crude Protein (Min) | 20.00% |
|---------------------|--------------|
| Crude Fat (Min) | 2.50% |
| Crude Fiber (Max) | 10.00% |
| Calcium (Min) | 3.50% |
| Calcium (Max) | 4.50% |
| Phosphorus (Min) | 0.75% |
| Salt (Min) | 15.00% |
| Salt (Max) | |
| Magnesium (Min) | 4.50% |
| Potassium (Min) | |
| Copper (Min) | |
| Manganese (Min) | |
| Selenium (Min) | |
| Zinc (Min) | |
| Vitamin A (Min) | 25,000 IU/lb |
| Vitamin D-3 (Min) | 5,000 IU/lb |
| Vitamin E (Min) | |
| | |

CAUTION: Use as directed. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

NET WEIGHT: 33.3 LB (15.12 KG)

INGREDIENTS

Processed Grain By-Products, Plant Protein Products, Salt, Magnesium Oxide, Molasses Products, Calcium Hydroxide, Monocalcium Phosphate, Dicalcium Phosphate, Calcium Carbonate, Hydrogenated Soybean Oil, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement and Mineral Oil.

INTRODUCTORY PERIOD

Following a 7 to 10 day acclimation period, average consumption should be 1 to 2 pounds for mature cattle with unrestricted forage supplies. Position the blocks 50 to 70 feet from the loafing, feeding and watering areas. Some repositioning of the blocks may be necessary as the animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption, further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of the blocks does not correct the situation, remove the blocks and evaluate the overall feeding program. DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.

Manufactured By: **Ridley Block Operations** Mankato, MN 56001 www.ultralvx.com 888-718-3493

For Beef Cattle on Pasture

FEEDING RECOMMENDATIONS

- Feed at a rate of one block per 5 to 15 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
- Place blocks where animals congregate. Locate blocks throughout the 2 pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to prevent crowding - 10 feet minimum.
- 3. Feed blocks continuously along with plentiful sources of forage and clean, fresh water. Cattle should not run out of blocks. When a block is two-thirds consumed, provide a new block near it. Place very small pieces in a feed pan or trough to be cleaned up.
- Cattle should consume 1 to 2 pounds per head daily based on a 1,000-lb 4 animal
- Consumption of this block may vary depending upon animal body 5. condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
- If additional mineral and vitamin fortification is desired, feed an appropriate 6. ULTRALYX® self-fed mineral supplement in addition to this supplement.

For additional information, please contact your ULTRALYX® representative.

Batch #



GUARANTEED ANALYSIS

| Crude Protein (Min) | 20.00% |
|---|---|
| Crude Fat (Min) | 2.50% |
| Crude Fiber (Max) | |
| Calcium (Min) | |
| Calcium (Max) | |
| Phosphorus (Min) | |
| Salt (Min) | |
| Salt (Max) | |
| Magnesium (Min) | |
| Potassium (Min) | |
| Copper (Min) | |
| | |
| Manganese (Min) | |
| Manganese (Min) Selenium (Min) | |
| Selenium (Min) | 3.3 ppm |
| Selenium (Min) Zinc (Min) | 3.3 ppm 300 ppm |
| Selenium (Min) Zinc (Min) Vitamin A (Min) | 3.3 ppm 300 ppm 25,000 IU/lb |
| Selenium (Min) Zinc (Min) | 3.3 ppm 300 ppm 25,000 IU/lb 5,000 IU/lb |

CAUTION: Use as directed. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

INGREDIENTS

Processed Grain By-Products, Plant Protein Products, Salt, Magnesium Oxide, Molasses Products, Calcium Hydroxide, Monocalcium Phosphate, Dicalcium Phosphate, Calcium 2. Carbonate, Hydrogenated Soybean Oil, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement and Mineral Oil

INTRODUCTORY PERIOD

Following a 7 to 10 day acclimation period, average consumption should be 1 to 2 pounds for mature cattle with unrestricted forage supplies. Position the blocks 50 to 70 feet from the loafing, feeding and watering areas. Some repositioning of the blocks may be necessary as the animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption, further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of the blocks does not correct the situation, remove the blocks and evaluate the overall feeding program. DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.

Manufactured By: **Ridley Block Operations** Mankato, MN 56001 www.ultralvx.com 888-718-3493

For Beef Cattle on Pasture

UltraPro[™] 20% AN Block

FEEDING RECOMMENDATIONS

- Feed at a rate of one block per 5 to 15 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
- Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to prevent crowding - 10 feet minimum.
- 3. Feed blocks continuously along with plentiful sources of forage and clean, fresh water. Cattle should not run out of blocks. When a block is two-thirds consumed, provide a new block near it. Place very small pieces in a feed pan or trough to be cleaned up.
- 4. Cattle should consume 1 to 2 pounds per head daily based on a 1,000-lb animal
- Consumption of this block may vary depending upon animal body 5. condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
- 6. If additional mineral and vitamin fortification is desired, feed an appropriate ULTRALYX® self-fed mineral supplement in addition to this supplement.

For additional information, please contact your ULTRALYX® representative.



