

# 16:8 Goat Mineral with Rumensin®

# **MEDICATED**

# For Further Manufacture of Feed

For the prevention of coccidiosis caused by Eimeria crandallis, Eimeria chistenseni and Eimeria ninakohlyakimovae in goats

#### **Active Drug Ingredient**

Monensin	960 g/ton
Guaranteed Analysis	
Calcium (Min)	
Calcium (Max)	17.40%
Phosphorus (Min)	8 00%
Salt (Min)	15.00%
Salt (Max)	18.00%
Magnesium (Min)	2.00%
Copper (Min)	1,500 ppm
Copper (Max)	
Manganese (Min)	
Selenium (Min)	50 ppm
Zinc (Min)	
Vitamin A (Min)	300,000 IU/lb
Vitamin D-3 (Min)	75,000 IU/lb
Vitamin E (Min)	400 IU/lb

#### Ingredients

Monocalcium Phosphate, Dicalcium Phosphate, Calcium Carbonate, Salt, Magnesium Oxide, Processed Grain By-Products, Molasses Products, Hydrogenated Soybean Oil, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil and Red Iron Oxide.

Feeding and Management
ULTRALYX® 16:8 Goat Mineral with Rumensin® is a medicated 2:1 calcium to phosphorus mixing mineral scientifically formulated to help prevent coccidiosis while providing for the unique trace mineral and vitamin needs of goats. All ingredients are selected for high biological availability and stability to deliver optimal nutrition to your goats.

Mixing Directions: Thoroughly mix 41.65 lbs. of ULTRALYX® 16:8 Goat Mineral with Rumensin® with 1,958.35 lbs. of non-medicated feed or mix 1.04 lbs. of ULTRALYX® 16:8 Goat Mineral with Rumensin® with 48.96 lbs of non-medicated feed to create a complete feed providing 20 g per ton of monensin. Feed as the sole ration to goats maintained in confinement. Provide clean, fresh water to all animals at all times.

**CAUTION: Use only as directed.** Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by equines has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats. Consumption of selenium should not exceed 0.7 mg per head daily. The addition to feed of higher levels of this premix containing selenium is not permitted.

WARNING: This product, which contains supplemental copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

For additional information, please contact your ULTRALYX® representative.

Rumensin® is a registered trademark of Elanco Animal Health, Indianapolis, IN

NET WT 25 LB (11.34 kg)

Manufactured By: **Ridley Block Operations** Mankato, MN 56001 www.ultralvx.com 1-888-718-3493

Batch #



# 16:8 Goat Mineral with Rumensin®

# **MEDICATED**

# For Further Manufacture of Feed

For the prevention of coccidiosis caused by Eimeria crandallis, Eimeria chistenseni and Eimeria ninakohlyakimovae in goats.

#### **Active Drug Ingredient**

Monensin	960 g/ton
Guaranteed Analysis	
Calcium (Min)	14.50%
Calcium (Max)	17.40%
Phosphorus (Min)	8.00%
Salt (Min)	15.00%
Salt (Max)	18.00%
Magnesium (Min)	2.00%
Copper (Min)	1,500 ppm
Copper (Max)	
Manganese (Min)	4,000 ppm
Selenium (Min)	50 ppm
Zinc (Min)	4,000 ppm
Vitamin A (Min)	300,000 IU/lb
Vitamin D-3 (Min)	75,000 IU/lb
Vitamin E (Min)	400 IU/lb

#### Ingredients

Monocalcium Phosphate, Dicalcium Phosphate, Calcium Carbonate, Salt, Magnesium Oxide, Processed Grain By-Products, Molasses Products, Hydrogenated Soybean Oil, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil and Red Iron Oxide.

Feeding and Management ULTRALYX® 16:8 Goat Mineral with Rumensin® is a medicated 2:1 calcium to phosphorus mixing mineral scientifically formulated to help prevent coccidiosis while providing for the unique trace mineral and vitamin needs of goats. All ingredients are selected for high biological availability and stability to deliver optimal nutrition to your goats.

Mixing Directions: Thoroughly mix 41.65 lbs. of ULTRALYX® 16:8 Goat Mineral with Rumensin® with 1,958.35 lbs. of non-medicated feed or mix 1.04 lbs. of ULTRALYX® 16:8 Goat Mineral with Rumensin® with 48.96 lbs of non-medicated feed to create a complete feed providing 20 g per ton of monensin. Feed as the sole ration to goats maintained in confinement. Provide clean, fresh water to all animals at all times.

CAUTION: Use only as directed. Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by equines has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats. Consumption of selenium should not exceed 0.7 mg per head daily. The addition to feed of higher levels of this premix containing selenium is not permitted.

WARNING: This product, which contains supplemental copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

For additional information, please contact your ULTRALYX® representative.

Rumensin® is a registered trademark of Elanco Animal Health, Indianapolis, IN

NET WT 25 LB (11.34 kg)

Batch #

Manufactured By: **Ridley Block Operations** Mankato, MN 56001 www.ultralyx.com 1-888-718-3493