# TRIPLE TRUST 20-15 RED

# **CALF MILK REPLACER**

Complete Milk Replacer to be fed to herd replacement calves.

## GUARANTEED ANALYSIS

Crude Protein	,	Min	 20.00	%
Crude Fat	,	Min	 15.00	8
Crude Fiber	,	Max	 0.50	8
Calcium	,	Min	 0.75	%
Calcium	,	Max	 1.25	%
Phosphorus	,	Min	 0.70	%
Vitamin A	,	Min	 35000	IU/lb.
Vitamin D3	,	Min	 7500	IU/lb.
Vitamin E	,	Min	 150	IU/lb.

#### INGREDIENTS

Dried whey, protein modified soy flour, animal fat (preserved with BHA & BHT), dried whey product, lecithin, calcium carbonate, hydrolyzed yeast, casein, L-lysine, DL-methionine, dicalcium phosphate, xanthan gum, calcium silicate, dried whey protein concentrate, vitamin E supplement, dried skimmed milk, ethoxylated mono and diglycerides, propylene glycol, choline chloride, ferrous sulfate, artificial flavor, zinc sulfate, vitamin A supplement, manganese sulfate, copper sulfate, ascorbic acid, vitamin Bl2 supplement, niacin supplement, biotin, vitamin Bl2 supplement, selenium yeast, brewer's dried yeast, calcium pantothenate, riboflavin supplement, menadione sodium bisulfite complex (source of vitamin K activity), pyridoxine hydrochloride, thiamine mononitrate, folic acid, ethylenediamine dihydriodide, cobalt sulfate

Important: Follow mixing and feeding directions.

Manufactured for: Central Farm Supply of KY 7500 Grade Lane Louisville, KY 40219

NET WEIGHT 50 LB (22.67 kg)

01-9282

### MIXING DIRECTIONS

Enclosed is a cup that holds approximately 12 ounces of dry milk replacer powder. Always weigh powder for accurate mixing.

Add 12 ounces (1 enclosed cup) of milk replacer powder to 2 quarts of warm water (110-115°F) and mix thoroughly. For large batches, mix 7.5 lbs. (10 full cups) of milk replacer powder to 5 gallons of warm water.

# FEEDING DIRECTIONS

Large breeds: Feed 2.3 quarts reconstituted milk replacer (100°F) twice daily.

Adjust amount fed based on calf body weight.

OFFER FRESH WATER AT ALL TIMES

# CALF CARE RECOMMENDATIONS

- 1)Feed large breed calves 4 quarts of high quality colostrum within 2 hours of birth and again 12 hours later. Adjust amounts for small breed calves. If colostrum quality or quantity is low, consider using a colostrum supplement or replacer.
- 2)Begin feeding milk replacer on day two.
  3)Begin feeding calf starter free-choice in
- 3) Begin feeding calf starter free-choice in the first week. Feed a small amount at first and replace daily.
- 4)Calves can be weaned from milk replacer successfully when intake of calf starter reaches 2 pounds daily for 3 consecutive days.
- 5)Environmental conditions may alter feeding rate. See your nutritionist for details.

The benefits of a comprehensive calf care program can result in more productive and profitable livestock.

3801 REV. 0317