

TRIPLE TRUST 20-20 NT1333 GOLD

CALF MILK REPLACER

Type C Medicated Feed

Complete Milk Replacer to be fed to herd replacement calves.

For calves and beef and non-lactating dairy cattle for treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia (shipping fever complex) caused by *Pasteurella multocida* susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by *E. coli* susceptible to neomycin.

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

ACTIVE DRUG INGREDIENT(S)
Neomycin Sulfate 1333 g/ton
Oxytetracycline 1333 g/ton

GUARANTEED ANALYSIS

Crude Protein	, Min	20.00 %
Crude Fat	, Min	20.00 %
Crude Fiber	, Max	0.15 %
Calcium	, Min	0.75 %
Calcium	, Max	1.25 %
Phosphorus	, Min	0.70 %
Vitamin A	, Min	35000 IU/lb.
Vitamin D3	, Min	7500 IU/lb.
Vitamin E	, Min	150 IU/lb.

INGREDIENTS
Dried whey product, dried whey protein concentrate, dried whey, animal fat (preserved with BHA & BHT), lecithin, calcium carbonate, citric acid (preservative), L-lysine, calcium silicate, dried skimmed milk, vitamin E supplement, DL-methionine, artificial flavor, iron proteinate, choline chloride, zinc proteinate, manganese proteinate, vitamin A supplement, copper proteinate, ascorbic acid, vitamin B12 supplement, niacin supplement, selenium yeast, brewer's dried yeast, ethoxylated mono and diglycerides, propylene glycol, biotin, vitamin D3 supplement, calcium pantothenate, riboflavin supplement, menadione sodium bisulfite complex (source of vitamin K activity), pyridoxine hydrochloride, thiamine mononitrate, folic acid, cobalt proteinate, ethylenediamine dihydriodide

Important: Follow mixing and feeding directions.

****CAUTION****

If symptoms persist after using for 2 or 3 days, consult a veterinarian. Treatment should continue 24 to 48 hours beyond remission of disease symptoms.

*** * * WARNING : * * ***

Withdraw from feed 5 days prior to slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

Manufactured for:
Central Farm Supply of KY
7500 Grade Lane
Louisville, KY 40219

NET WEIGHT 50 LB (22.67 kg)

MIXING DIRECTIONS

Using the chart as a guide, mix the appropriate amount of milk replacer powder with clean, warm water (110°F), as recommended below per calf bodyweight. Feed the milk replacer solution immediately at a feeding temperature of 100°F.

The enclosed cup holds approximately 12 ounces of powder. For accurate feeding, weigh powder.

FEEDING DIRECTIONS

Feed to provide each calf 10 mg. of neomycin sulfate and 10 mg. oxytetracycline per pound of bodyweight daily.

Feed continuously for 7 to 14 days in place of normal milk replacer.

Feed the medicated milk replacer solution TWICE daily according to the following schedule.

CALF WT. (lbs.)	NEOMYCIN & OXYTETRACYCLINE (mg/day)	POWDER (oz.)	WATER (quarts)	FEEDINGS PER DAY
100	1000	12	2	2
125	1250	15	2.5	2
150	1500	18	3	2

OFFER FRESH WATER AT ALL TIMES

CALF CARE RECOMMENDATIONS

- 1) Feed large breed calves 4 quarts of high quality colostrum within 2 hours of birth and again 12 hours later. Adjust amounts for small breed calves.
- 2) Begin feeding milk replacer on day two.
- 3) Begin feeding calf starter free-choice in the first week. Feed a small amount at first and replace daily.
- 4) Calves can be weaned from milk replacer successfully when intake of calf starter reaches 2 pounds daily for 3 consecutive days.
- 5) Environmental conditions may alter feeding rate. See your nutritionist for details.

The benefits of a comprehensive calf care program can result in more productive and profitable livestock.