



S-G-D

Low Moisture Block

For Sheep, Goats and Deer on Pasture

Guaranteed Analysis:

Crude Protein (Min)	15.00%
Crude Fat (Min)	5.00%
Crude Fiber (Max)	2.50%
Acid Detergent Fiber (Max)	8.00%
Calcium (Min)	1.30%
Calcium (Max)	1.80%
Phosphorus (Min)	0.90%
Magnesium (Min)	0.40%
Potassium (Min)	4.00%
Copper (Min)	5 ppm
Copper (Max)	35 ppm
Manganese (Min)	400 ppm
Selenium (Min)	3.5 ppm
Zinc (Min)	1,600 ppm
Vitamin A (Min)	55,000 IU/lb
Vitamin D-3 (Min)	5,500 IU/lb
Vitamin E (Min)	425 IU/lb

CAUTION: Use as directed. Consumption of selenium should not exceed 0.7 mg per head daily.

Ingredients:

Molasses Products, Processed Grain By-Products, Monocalcium Phosphate, Dicalcium Phosphate, Hydrolyzed Feather Meal, Hydrolyzed Vegetable Oil, Calcium Carbonate, Potassium Chloride, Magnesium Oxide, Ferrous Sulfate, Manganous Oxide, Zinc Oxide, Manganese Sulfate, Zinc Sulfate, Sodium Selenite, Mineral Oil, Calcium Iodate, Cobalt Sulfate, Vitamin A Supplement, Vitamin D-3 Supplement and Vitamin E Supplement.

INTRODUCTORY PERIOD:

Following a 7 to 10 day acclimation period, average consumption should be 1 to 3 ounces for sheep, goats and deer with unrestricted forage supplies. Position the blocks 50 to 75 feet from the loafing, feeding and watering areas during the first week of feeding. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If over consumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

FEEDING RECOMMENDATIONS

1. Provide 1 block for every 10 to 20 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to minimize crowding - 10 feet minimum.
3. Feed blocks continuously along with plentiful sources of forage and clean, fresh water. Animals should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
4. Animals should consume 1 to 3 ounces per head per day.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional vitamin and mineral fortification is desired, provide an appropriate Triple Trust® self-fed supplement in addition to this supplement. **This product does not contain salt; provide a source of free choice salt.**

For more information, contact your Triple Trust® representative.

NET WEIGHT: 50 LBS. (22.7 KG)

Manufactured By:
Ridley Block Operations
Mankato, MN 56001

Batch #

25160
0116



S-G-D

Low Moisture Block

For Sheep, Goats and Deer on Pasture

Guaranteed Analysis:

Crude Protein (Min)	15.00%
Crude Fat (Min)	5.00%
Crude Fiber (Max)	2.50%
Acid Detergent Fiber (Max)	8.00%
Calcium (Min)	1.30%
Calcium (Max)	1.80%
Phosphorus (Min)	0.90%
Magnesium (Min)	0.40%
Potassium (Min)	4.00%
Copper (Min)	5 ppm
Copper (Max)	35 ppm
Manganese (Min)	400 ppm
Selenium (Min)	3.5 ppm
Zinc (Min)	1,600 ppm
Vitamin A (Min)	55,000 IU/lb
Vitamin D-3 (Min)	5,500 IU/lb
Vitamin E (Min)	425 IU/lb

CAUTION: Use as directed. Consumption of selenium should not exceed 0.7 mg per head daily.

Ingredients:

Molasses Products, Processed Grain By-Products, Monocalcium Phosphate, Dicalcium Phosphate, Hydrolyzed Feather Meal, Hydrolyzed Vegetable Oil, Calcium Carbonate, Potassium Chloride, Magnesium Oxide, Ferrous Sulfate, Manganous Oxide, Zinc Oxide, Manganese Sulfate, Zinc Sulfate, Sodium Selenite, Mineral Oil, Calcium Iodate, Cobalt Sulfate, Vitamin A Supplement, Vitamin D-3 Supplement and Vitamin E Supplement.

INTRODUCTORY PERIOD:

Following a 7 to 10 day acclimation period, average consumption should be 1 to 3 ounces for sheep, goats and deer with unrestricted forage supplies. Position the blocks 50 to 75 feet from the loafing, feeding and watering areas during the first week of feeding. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If over consumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

FEEDING RECOMMENDATIONS

1. Provide 1 block for every 10 to 20 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to minimize crowding - 10 feet minimum.
3. Feed blocks continuously along with plentiful sources of forage and clean, fresh water. Animals should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
4. Animals should consume 1 to 3 ounces per head per day.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional vitamin and mineral fortification is desired, provide an appropriate Triple Trust® self-fed supplement in addition to this supplement. **This product does not contain salt; provide a source of free choice salt.**

For more information, contact your Triple Trust® representative.

NET WEIGHT: 50 LBS. (22.7 KG)

Manufactured By:
Ridley Block Operations
Mankato, MN 56001

Batch #

25160
0116