



20-1 Dry Cow

Low Moisture Block

For Dairy Cattle on Pasture

Guaranteed Analysis:

Crude Protein (Min).....	20.00%
(This includes not more than 8.00% protein equivalent from non-protein nitrogen.)	
Crude Fat (Min).....	2.00%
Crude Fiber (Max).....	2.00%
Acid Detergent Fiber (Max).....	8.00%
Calcium (Min).....	2.50%
Calcium (Max).....	3.50%
Phosphorus (Min).....	3.00%
Magnesium (Min).....	0.70%
Potassium (Min).....	2.00%
Cobalt (Min).....	10 ppm
Copper (Min).....	500 ppm
Iron (Min).....	1,850 ppm
Manganese (Min).....	1,500 ppm
Selenium (Min).....	6.6 ppm
Zinc (Min).....	1,500 ppm
Vitamin A (Min).....	100,000 IU/lb
Vitamin D-3 (Min).....	30,000 IU/lb
Vitamin E (Min).....	1,000 IU/lb

CAUTION: Use as directed. Feeding added selenium in excess of 0.3 ppm in the total diet is prohibited.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

Ingredients:

Molasses Products, Processed Grain By-Products, Monocalcium Phosphate, Dicalcium Phosphate, Hydrolyzed Feather Meal, Urea, Magnesium Oxide, Hydrolyzed Vegetable Oil, Calcium Carbonate, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Ferrous Sulfate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement and Mineral Oil.

INTRODUCTORY PERIOD:

Following a 7 to 10 day acclimation period, average consumption should be 0.5 to 1 pounds for mature cattle with unrestricted forage supplies. Position the blocks 50 to 75 feet from the loafing, feeding and watering areas during the first week of feeding. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If over consumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

NET WEIGHT: 200 LBS. (90.7 KG)

Manufactured By:
Ridley Block Operations
Mankato, MN 56001

Batch #

25166
0116

FEEDING RECOMMENDATIONS

1. Feed at a rate of one block per 15 to 30 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to prevent crowding - 20-30 feet minimum.
3. Feed blocks continuously along with plentiful sources of forage and clean, fresh water. Animals should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
4. Cattle should consume 0.5 to 1 pounds per head daily based on a 1,000-pound animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional vitamin and mineral fortification is desired, provide an appropriate Triple Trust® self-fed supplement in addition to this supplement. **This product does not contain salt, provide a source of free choice salt. Limit salt intake to 1 ounce per head daily.**

For more information contact your Triple Trust® representative.



20-1 Dry Cow

Low Moisture Block

For Dairy Cattle on Pasture

Guaranteed Analysis:

Crude Protein (Min).....	20.00%
(This includes not more than 8.00% protein equivalent from non-protein nitrogen.)	
Crude Fat (Min).....	2.00%
Crude Fiber (Max).....	2.00%
Acid Detergent Fiber (Max).....	8.00%
Calcium (Min).....	2.50%
Calcium (Max).....	3.50%
Phosphorus (Min).....	3.00%
Magnesium (Min).....	0.70%
Potassium (Min).....	2.00%
Cobalt (Min).....	10 ppm
Copper (Min).....	500 ppm
Iron (Min).....	1,850 ppm
Manganese (Min).....	1,500 ppm
Selenium (Min).....	6.6 ppm
Zinc (Min).....	1,500 ppm
Vitamin A (Min).....	100,000 IU/lb
Vitamin D-3 (Min).....	30,000 IU/lb
Vitamin E (Min).....	1,000 IU/lb

CAUTION: Use as directed. Feeding added selenium in excess of 0.3 ppm in the total diet is prohibited.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

Ingredients:

Molasses Products, Processed Grain By-Products, Monocalcium Phosphate, Dicalcium Phosphate, Hydrolyzed Feather Meal, Urea, Magnesium Oxide, Hydrolyzed Vegetable Oil, Calcium Carbonate, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Ferrous Sulfate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement and Mineral Oil.

INTRODUCTORY PERIOD:

Following a 7 to 10 day acclimation period, average consumption should be 0.5 to 1 pounds for mature cattle with unrestricted forage supplies. Position the blocks 50 to 75 feet from the loafing, feeding and watering areas during the first week of feeding. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If over consumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

NET WEIGHT: 200 LBS. (90.7 KG)

Manufactured By:
Ridley Block Operations
Mankato, MN 56001

Batch #

25166
0116

FEEDING RECOMMENDATIONS

1. Feed at a rate of one block per 15 to 30 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to prevent crowding - 20-30 feet minimum.
3. Feed blocks continuously along with plentiful sources of forage and clean, fresh water. Animals should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
4. Cattle should consume 0.5 to 1 pounds per head daily based on a 1,000-pound animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional vitamin and mineral fortification is desired, provide an appropriate Triple Trust® self-fed supplement in addition to this supplement. **This product does not contain salt, provide a source of free choice salt. Limit salt intake to 1 ounce per head daily.**

For more information contact your Triple Trust® representative.