

# BC Calving and Rebreeding Tub

*Compressed Block*

For Beef Cattle on Pasture

## Guaranteed Analysis:

Crude Protein (Min)	20.00%
Crude Fat (Min)	5.50%
Crude Fiber (Max)	3.75%
Calcium (Min)	4.50%
Calcium (Max)	5.50%
Phosphorus (Min)	1.40%
Salt (Min)	4.80%
Salt (Max)	5.80%
Magnesium (Min)	4.00%
Potassium (Min)	2.30%
Cobalt (Min)	11 ppm
Copper (Min)	400 ppm
Iodine (Min)	30 ppm
Manganese (Min)	1,200 ppm
Selenium (Min)	6.6 ppm
Zinc (Min)	1,600 ppm
Vitamin A (Min)	50,000 IU/lb
Vitamin D-3 (Min)	11,000 IU/lb
Vitamin E (Min)	80 IU/lb

**CAUTION: Use as directed.** Consumption of selenium should not exceed 3 mg per head daily.

Manufactured By:  
**Ridley Block Operations**  
Mankato, MN 56001

## Ingredients:

Plant Protein Products, Processed Grain By-Products, Magnesium Oxide, Grain Products, Calcium Hydroxide, Salt, Monocalcium Phosphate, Dicalcium Phosphate, Potassium Chloride, Molasses Products, Hydrolyzed Vegetable Oil, Calcium Carbonate, Manganese Amino Acid Chelate, Manganous Oxide, Manganese Sulfate, Zinc Amino Acid Chelate, Zinc Oxide, Zinc Sulfate, Copper Amino Acid Chelate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement and Mineral Oil.

**WARNING:** This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

**Introductory Period:** Following a 7 to 10 day acclimation period, average consumption should be 0.5 to 1 pound per head per day for mature cattle with unrestricted forage supplies. Position the blocks 50 to 75 feet from loafing, feeding and watering areas. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

## Feeding Recommendations:

1. Provide 1 block per 15 to 30 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing, feeding and watering areas. Adequate distance must be maintained between blocks to minimize crowding – 10 feet minimum.
3. Feed blocks continuously along with a plentiful source of forage and clean, fresh water. Cattle should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
4. Cattle should consume an average of 0.5 to 1 pound per head daily based on a 1,000-pound animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional mineral and vitamin fortification is desired, provide an appropriate free choice mineral supplement in addition to this supplement.

For additional information, contact your Ridley Block Operations representative.

**Net Weight: 200 LBS. (90.7 KG)**

Batch #

**27081**

0616

# BC Calving and Rebreeding Tub

*Compressed Block*

For Beef Cattle on Pasture

## Guaranteed Analysis:

Crude Protein (Min)	20.00%
Crude Fat (Min)	5.50%
Crude Fiber (Max)	3.75%
Calcium (Min)	4.50%
Calcium (Max)	5.50%
Phosphorus (Min)	1.40%
Salt (Min)	4.80%
Salt (Max)	5.80%
Magnesium (Min)	4.00%
Potassium (Min)	2.30%
Cobalt (Min)	11 ppm
Copper (Min)	400 ppm
Iodine (Min)	30 ppm
Manganese (Min)	1,200 ppm
Selenium (Min)	6.6 ppm
Zinc (Min)	1,600 ppm
Vitamin A (Min)	50,000 IU/lb
Vitamin D-3 (Min)	11,000 IU/lb
Vitamin E (Min)	80 IU/lb

**CAUTION: Use as directed.** Consumption of selenium should not exceed 3 mg per head daily.

Manufactured By:  
**Ridley Block Operations**  
Mankato, MN 56001

## Ingredients:

Plant Protein Products, Processed Grain By-Products, Magnesium Oxide, Grain Products, Calcium Hydroxide, Salt, Monocalcium Phosphate, Dicalcium Phosphate, Potassium Chloride, Molasses Products, Hydrolyzed Vegetable Oil, Calcium Carbonate, Manganese Amino Acid Chelate, Manganous Oxide, Manganese Sulfate, Zinc Amino Acid Chelate, Zinc Oxide, Zinc Sulfate, Copper Amino Acid Chelate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement and Mineral Oil.

**WARNING:** This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

**Introductory Period:** Following a 7 to 10 day acclimation period, average consumption should be 0.5 to 1 pound per head per day for mature cattle with unrestricted forage supplies. Position the blocks 50 to 75 feet from loafing, feeding and watering areas. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

## Feeding Recommendations:

1. Provide 1 block per 15 to 30 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing, feeding and watering areas. Adequate distance must be maintained between blocks to minimize crowding – 10 feet minimum.
3. Feed blocks continuously along with a plentiful source of forage and clean, fresh water. Cattle should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
4. Cattle should consume an average of 0.5 to 1 pound per head daily based on a 1,000-pound animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional mineral and vitamin fortification is desired, provide an appropriate free choice mineral supplement in addition to this supplement.

For additional information, contact your Ridley Block Operations representative.

**Net Weight: 200 LBS. (90.7 KG)**

Batch #

**27081**

0616