



# 16% Goat

For All Classes of Goats

## GUARANTEED ANALYSIS

Crude Protein (Min) .....	16.00%
Crude Fat (Min) .....	3.00%
Crude Fiber (Max) .....	5.00%
Acid Detergent Fiber (Max) .....	7.00%
Calcium (Min) .....	6.00%
Calcium (Max) .....	7.20%
Phosphorus (Min) .....	2.00%
Salt (Min) .....	10.00%
Salt (Max) .....	12.00%
Magnesium (Min) .....	2.00%
Copper (Min) .....	200 ppm
Copper (Max) .....	250 ppm
Manganese (Min) .....	2,000 ppm
Selenium (Min) .....	6.0 ppm
Zinc (Min) .....	2,000 ppm
Vitamin A (Min) .....	100,000 IU/lb
Vitamin D-3 (Min) .....	10,000 IU/lb
Vitamin E (Min) .....	100 IU/lb

**CAUTION:** Use as directed. Consumption of selenium should not exceed 0.7 mg per head daily.

**WARNING:** This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

**NET WEIGHT: 25 LBS (11.34 KG)**

## INGREDIENTS

Processed Grain By-Products, Calcium Carbonate, Salt, Molasses Products, Monocalcium Phosphate, Dicalcium Phosphate, Plant Protein Products, Magnesium Oxide, Calcium Hydroxide, Manganous Oxide, Manganese Sulfate, Manganese Proteinates, Zinc Oxide, Zinc Sulfate, Zinc Proteinates, Copper Sulfate, Copper Proteinates, Sodium Selenite, Selenium Yeast, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Cobalt Proteinates, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil.

## INTRODUCTORY PERIOD

Following a 7 to 10 day acclimation period, average consumption should be 2 to 4 ounces per head daily for goats with unrestricted forage supplies. Position the blocks 50 to 70 feet from the loafing, feeding and watering areas. Some repositioning of the blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption, further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of the blocks does not correct the situation, remove the blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

## FEEDING RECOMMENDATIONS

1. Feed 1 block per 5 to 10 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place the blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing, feeding and watering areas. Adequate distance must be maintained between blocks to minimize crowding - 10 feet minimum.
3. Feed blocks continuously along with a plentiful source of forage and clean, fresh water. Animals should not run out of blocks. When a block is two-thirds consumed, provide a new block near it. Place very small pieces in a feed trough to be cleaned up.
4. Goats should consume an average of 2 to 4 ounces per head daily.
5. Consumption may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional mineral and vitamin fortification is desired, provide an appropriate Triple Trust® self-fed mineral supplement in addition to this supplement.

For additional information, please contact your Triple Trust® representative.

Manufactured By:  
Ridley Block Operations  
Mankato, MN 56001



**56236**

0916