



20% Equine

For All Classes of Equines

GUARANTEED ANALYSIS

Crude Protein (Min)	20.00%
Crude Fat (Min)	2.00%
Crude Fiber (Max)	6.50%
Acid Detergent Fiber (Max)	6.00%
Neutral Detergent Fiber (Max)	18.00%
Calcium (Min)	3.00%
Calcium (Max)	4.00%
Phosphorus (Min)	2.00%
Salt (Min)	19.00%
Salt (Max)	22.80%
Copper (Min).....	125 ppm
Manganese (Min).....	270 ppm
Selenium (Min)	3.3 ppm
Zinc (Min).....	300 ppm
Vitamin A (Min)	25,000 IU/lb
Vitamin D-3 (Min).....	2,500 IU/lb
Vitamin E (Min)	25 IU/lb

CAUTION: Use as directed. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

NET WEIGHT: 25 LBS (11.34 KG)

INGREDIENTS

Processed Grain By-Products, Salt, Plant Protein Products, Molasses Products, Monocalcium Phosphate, Dicalcium Phosphate, Calcium Hydroxide, Calcium Carbonate, *Saccharomyces cerevisiae* Yeast Culture, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Selenium Yeast, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil.

INTRODUCTORY PERIOD

Following a 7 to 10 day acclimation period, average consumption should be 1 to 2 pounds per head per day for mature horses with unrestricted forage supplies. Position the blocks 50 to 75 feet from loafing, feeding and watering areas. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of the blocks does not correct the situation, remove the blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED HORSES FREE ACCESS TO THIS BLOCK.**

FEEDING RECOMMENDATIONS

1. Feed one block per 5 to 15 horses. All horses should have equal access to blocks. Timid and smaller horses need their share.
2. Place blocks where horses congregate. Locate blocks throughout the pasture at loafing, grazing, feeding and watering areas. Adequate distance must be maintained between blocks to minimize crowding - 10 feet minimum.
3. Feed blocks continuously along with plentiful sources of forage and clean, fresh water. Horses should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
4. Horses should consume an average of 1 to 2 pounds per head daily based on a 1,000-pound animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages fed, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional mineral and vitamin fortification is desired, provide an appropriate Triple Trust® self-fed mineral supplement in addition to this supplement.

For additional information, please contact your Triple Trust® representative.

Manufactured By:
Ridley Block Operations
Mankato, MN 56001



56237

0916