



Stress and Weaning

Low Moisture Block

For Beef Cattle on Pasture

Guaranteed Analysis:

Crude Protein (Min)	16.00%
Crude Fat (Min)	5.00%
Crude Fiber (Max)	2.00%
Calcium (Min)	2.50%
Calcium (Max)	3.50%
Phosphorus (Min)	2.00%
Magnesium (Min)	0.70%
Potassium (Min)	4.00%
Cobalt (Min)	4.5 ppm
Copper (Min)	400 ppm
Iodine (Min)	9 ppm
Manganese (Min)	1,300 ppm
Selenium (Min)	4.4 ppm
Zinc (Min)	1,400 ppm
Vitamin A (Min)	200,000 IU/lb
Vitamin D-3 (Min)	20,000 IU/lb
Vitamin E (Min)	800 IU/lb
Thiamine (Min)	95 mg/lb

CAUTION: Use as directed. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

Ingredients:

Molasses Products, Processed Grain By-Products, Hydrolyzed Feather Meal, Monocalcium Phosphate, Dicalcium Phosphate, Hydrolyzed Vegetable Oil, Calcium Carbonate, Potassium Chloride, Magnesium Oxide, Manganese Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydrochloride, Calcium Iodate, Cobalt Carbonate, Thiamine Mononitrate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement and Mineral Oil.

INTRODUCTORY PERIOD:

Following a 7 to 10 day acclimation period, average consumption should be 1 to 1.5 pounds for mature cattle with unrestricted forage supplies. Position the blocks 50 to 75 feet from the loafing, feeding and watering areas during the first week of feeding. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If over consumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

FEEDING RECOMMENDATIONS

1. Feed at a rate of one block per 15 to 30 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to prevent crowding - 20-30 feet minimum.
3. Feed blocks continuously along with plentiful sources of forage and clean, fresh water. Animals should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
4. Cattle should consume 1 to 1.5 pounds per head daily based on a 1,000-pound animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional vitamin and mineral fortification is desired, provide an appropriate Triple Trust® self-fed supplement in addition to this supplement. **This product does not contain salt, provide a source of free choice salt.**

For more information contact your Triple Trust® representative.

Manufactured By:
Ridley Block Operations
Mankato, MN 56001

Batch #

25165

0116

NET WEIGHT: 200 LBS. (90.7 KG)