

**Guaranteed Analysis:**

Crude Protein (Min)	12.00%
Crude Fat (Min)	4.00%
Crude Fiber (Max)	2.50%
Calcium (Min)	1.00%
Calcium (Max)	1.50%
Phosphorus (Min)	1.00%
Salt (Min)	12.00%
Salt (Max)	14.40%
Magnesium (Min)	0.70%
Potassium (Min)	2.50%
Cobalt (Min)	4 ppm
Copper (Min)	220 ppm
Iodine (Min)	6 ppm
Manganese (Min)	750 ppm
Selenium (Min)	0.7 ppm
Zinc (Min)	750 ppm
Vitamin A (Min)	30,000 IU/lb
Vitamin D-3 (Min)	5,000 IU/lb
Vitamin E (Min)	220 IU/lb
Biotin (Min)	10 mg/lb
Thiamine (Min)	30 mg/lb

CAUTION: Use as directed. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

Horse Lix

Low Moisture Block

For Horses on Pasture

Ingredients:

Molasses Products, Processed Grain By-Products, Salt, Plant Protein Products, Monocalcium Phosphate, Dicalcium Phosphate, Magnesium Oxide, Hydrolyzed Vegetable Oil, Calcium Carbonate, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Biotin Supplement, Thiamine Mononitrate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement and Mineral Oil.

INTRODUCTORY PERIOD:

Following a 7 to 10 day acclimation period, average consumption should be 1 to 3 pounds for horses with unrestricted forage supplies. Position the blocks 50 to 75 feet from the loafing, feeding and watering areas during the first week of feeding. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If over consumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

FEEDING RECOMMENDATIONS

1. Provide 1 block for every 5 to 10 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to minimize crowding - 10 feet minimum.
3. Feed blocks continuously along with plentiful sources of forage and clean, fresh water. Horses should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
4. Horses should consume 1 to 3 pounds per head per day based on a 1,000-pound animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional vitamin and mineral fortification is desired, provide an appropriate Triple Trust® self-fed supplement in addition to this supplement.

For more information, contact your Triple Trust® representative.

NET WEIGHT: 200 LBS. (90.7 KG)

Manufactured By:
Ridley Block Operations
Mankato, MN 56001

Batch #

18772
0116