



All Purpose 12%

Low Moisture Block

For Cattle, Sheep and Goats on Pasture

Guaranteed Analysis:

Crude Protein (Min)	12.00%
Crude Fat (Min)	1.00%
Crude Fiber (Max)	2.50%
Calcium (Min)	1.00%
Calcium (Max)	1.50%
Phosphorus (Min)	0.80%
Potassium (Min)	2.00%
Cobalt (Min)	3 ppm
Copper (Min)	5 ppm
Copper (Max)	35 ppm
Iodine (Min)	12 ppm
Manganese (Min)	320 ppm
Selenium (Min)	4.4 ppm
Zinc (Min)	720 ppm
Vitamin A (Min)	80,000 IU/lb
Vitamin D-3 (Min)	8,000 IU/lb
Vitamin E (Min)	80 IU/lb

This product has been formulated for use in situations where cattle, sheep and goats are grazed together, and may not provide required nutrients for all these species in all situations.

CAUTION: Use as directed. Consumption of selenium should not exceed 3 mg per head per day for cattle and not exceed 0.7 mg per head per day for sheep or goats.

NET WEIGHT: 250 LBS. (113.4 KG)

Ingredients:

Molasses Products, Processed Grain By-Products, Monocalcium Phosphate, Dicalcium Phosphate, Calcium Carbonate, Ferrous Sulfate, Manganous Oxide, Zinc Oxide, Manganese Sulfate, Sodium Selenite, Zinc Sulfate, Mineral Oil, Calcium Iodate, Cobalt Sulfate, Sodium Molybdate, Ethylenediamine Dihydrochloride, Vitamin A Supplement, Vitamin D-3 Supplement and Vitamin E Supplement.

INTRODUCTORY PERIOD:

Following a 7 to 10 day acclimation period, average consumption should be 0.5 to 1 pound for mature cattle, and sheep and goats average consumption should be 1 to 3 ounces per head per day with unrestricted forage supplies. Position the blocks 50 to 75 feet from the loafing, feeding and watering areas during the first week of feeding. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If over consumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

FEEDING RECOMMENDATIONS

1. Provide 1 block for every 20 to 30 head of cattle and 10 to 20 head of sheep or goats. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to minimize crowding - 10 feet minimum. Feed blocks continuously along with plentiful sources of forage and clean, fresh water. Animals should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
3. Cattle should consume 0.5 to 1 pound per head daily based on a 1,000-pound animal. Sheep and goats should consume an average of 1 to 3 ounces per head daily.
4. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
5. If additional vitamin and mineral fortification is desired, provide an appropriate Triple Trust® self-fed supplement in addition to this supplement. **This product does not contain salt, provide a source of free choice salt.**

For more information, contact your Triple Trust® representative.

Manufactured By:
Ridley Block Operations
Mankato, MN 56001

Batch #

18758

1115