



Cattle Mag BioPlex® Block

For Beef Cattle on Pasture

GUARANTEED ANALYSIS

| | |
|-------------------------|---------------|
| Calcium (Min) | 3.00% |
| Calcium (Max) | 4.00% |
| Phosphorus (Min) | 0.40% |
| Salt (Min) | 12.00% |
| Salt (Max) | 14.40% |
| Magnesium (Min) | 14.00% |
| Copper (Min) | 440 ppm |
| Manganese (Min) | 1,800 ppm |
| Selenium (Min) | .26 ppm |
| Zinc (Min) | 2,100 ppm |
| Vitamin A (Min) | 100,000 IU/lb |
| Vitamin D-3 (Min) | 10,000 IU/lb |
| Vitamin E (Min) | 100 IU/lb |

CAUTION: Use as directed.

Consumption of selenium should not exceed 3 mg per head daily.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

BioPlex® is a registered trademark of Alltech.

INGREDIENTS

Processed Grain By-Products, Magnesium Oxide, Molasses Products, Salt, Calcium Carbonate, Calcium Hydroxide, Monocalcium Phosphate, Dicalcium Phosphate, Manganous Oxide, Manganese Sulfate, Manganese Proteinate, Zinc Oxide, Zinc Sulfate, Zinc Proteinate, Copper Sulfate, Copper Proteinate, Sodium Selenite, Selenium Yeast, Brewers Dried Yeast, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Cobalt Proteinate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement and Mineral Oil.

INTRODUCTORY PERIOD

Position the blocks 50 to 70 feet from the loafing, feeding and watering areas. Some repositioning of the blocks may be necessary as the animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption, further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of the blocks does not correct the situation, remove the blocks and evaluate the overall feeding program.

FEEDING RECOMMENDATIONS

1. Feed at a rate of one block per 5 to 15 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to prevent crowding - 10 feet minimum.
3. Feed blocks continuously along with plentiful sources of forage and clean, fresh water. Cattle should not run out of blocks. When a block is two-thirds consumed, provide a new block near it. Place very small pieces in a feed pan or trough to be cleaned up.
4. Cattle should consume 2 to 4 ounces per head daily based on a 1,000-lb animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. Feed as the sole source of salt, minerals or vitamins.

NET WEIGHT: 33.3 LB (15.12 KG)

For additional information, please contact your Triple Trust® representative.



Manufactured By: **Ridley Block Operations** • Mankato, MN 56001

10339

0816

