



Free Choice Rumensin® 1620 Mineral

TYPE C MEDICATED

A Free Choice Mineral for Growing Cattle

For increased rate of weight gain and for prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) 1,620 g/ton
(each lb. contains 810 mg)

GUARANTEED ANALYSIS

Calcium (Min).....9.0%
Calcium (Max)..... 10.5%
Phosphorus (Min).....6.0%
Salt (Min).....22.0%
Salt (Max).....26.0%
Potassium (Min).....1.0%
Copper (Min).....500 ppm
Iodine (min)..... 20 ppm
Manganese (Min).....2,100 ppm
Selenium (Min)..... 26 ppm
Zinc (Min).....1,560 ppm
Vitamin A (Min)..... 100,000 IU/lb
Vitamin D-3 (Min)..... 10,000 IU/lb
Vitamin E (Min).....10 IU/lb

INGREDIENTS

Monocalcium Phosphate, Dicalcium Phosphate, Salt, Roughage Products, Molasses Products, Calcium Carbonate, Processed Grain By-Products, Hydrogenated Soybean Oil, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Chloride, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Red Iron Oxide and Artificial Flavor.

SEE REVERSE SIDE FOR FEEDING DIRECTIONS AND PRECAUTIONARY STATEMENTS

Manufactured by
Ridley Block Operations
Mankato, MN 56001
www.ultralyx.com

NET WT 50 LB (22.7 kg)

Rumensin® is a registered trademark of Elanco Animal Health, Indianapolis, IN.

Batch #

10507
UL2304
10/13



Free Choice Rumensin® 1620 Mineral

TYPE C MEDICATED

A Free Choice Mineral for Growing Cattle

For increased rate of weight gain and for prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) 1,620 g/ton
(each lb. contains 810 mg)

GUARANTEED ANALYSIS

Calcium (Min) 9.0%
Calcium (Max) 10.5%
Phosphorus (Min).....6.0%
Salt (Min).....22.0%
Salt (Max).....26.0%
Potassium (Min).....1.0%
Copper (Min) 500 ppm
Iodine (min) 20 ppm
Manganese (Min) 2,100 ppm
Selenium (Min) 26 ppm
Zinc (Min)..... 1,560 ppm
Vitamin A (Min).....100,000 IU/lb
Vitamin D-3 (Min)10,000 IU/lb
Vitamin E (Min)..... 10 IU/lb

INGREDIENTS

Monocalcium Phosphate, Dicalcium Phosphate, Salt, Roughage Products, Molasses Products, Calcium Carbonate, Processed Grain By-Products, Hydrogenated Soybean Oil, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Chloride, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Red Iron Oxide and Artificial Flavor.

SEE REVERSE SIDE FOR FEEDING DIRECTIONS AND PRECAUTIONARY STATEMENTS

Manufactured by
Ridley Block Operations
Mankato, MN 56001
www.ultralyx.com

NET WT 50 LB (22.7 kg)

Rumensin® is a registered trademark of Elanco Animal Health, Indianapolis, IN.

Batch #

10507
UL2304
10/13

INSTRUCTIONS FOR FEEDING

For free choice feeding to growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers). Feed continuously on a free choice basis at the rate of 50 to 200 mg per head per day. Place in covered mineral feeders located near the animals water supply and/or loafing areas. Provide one feeder for each 20 head. Fill feeders with a quantity of **ULTRALYX® Free Choice Rumensin® 1620 Mineral** that will be consumed in seven days. It is essential to offer enough feeding stations to ensure that all animals have free access at all times to **ULTRALYX® Free Choice Rumensin® 1620 Mineral**. Cattle should consume 1 to 4 oz. per head daily which will provide the approved effective intake of 50 to 200 mg monensin. **NOTE:** Adequate consumption for cattle is dependent on good pasture conditions.

Provide fresh, clean water at all times. For additional information contact your ULTRALYX® representative.

CAUTION: During the first 5 days of feeding, cattle should receive no more than 100 mg/head/day. Do not feed additional salt or minerals. Do not mix with grain or other feeds. Stressed and/or water deprived cattle should be adapted to the pasture and to un-medicated mineral supplement before using **ULTRALYX® Free Choice Rumensin® 1620 Mineral**. Do not allow horses or other equines access to feeds containing monensin. Ingestions of monensin by horses has been fatal. Monensin medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions as reduced average daily gains may result. Do not feed to lactating goats. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: A withdrawal period has not been established in pre-ruminating calves. Do not use in calves to be processed for veal.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

Manufactured by
Ridley Block Operations
Mankato, MN 56001
www.ultralyx.com

NET WT 50 LB (22.7 kg)

Batch #

10507
UL2304
10/13

INSTRUCTIONS FOR FEEDING

For free choice feeding to growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers). Feed continuously on a free choice basis at the rate of 50 to 200 mg per head per day. Place in covered mineral feeders located near the animals water supply and/or loafing areas. Provide one feeder for each 20 head. Fill feeders with a quantity of **ULTRALYX® Free Choice Rumensin® 1620 Mineral** that will be consumed in seven days. It is essential to offer enough feeding stations to ensure that all animals have free access at all times to **ULTRALYX® Free Choice Rumensin® 1620 Mineral**. Cattle should consume 1 to 4 oz. per head daily which will provide the approved effective intake of 50 to 200 mg monensin. **NOTE:** Adequate consumption for cattle is dependent on good pasture conditions.

Provide fresh, clean water at all times. For additional information contact your ULTRALYX® representative.

CAUTION: During the first 5 days of feeding, cattle should receive no more than 100 mg/head/day. Do not feed additional salt or minerals. Do not mix with grain or other feeds. Stressed and/or water deprived cattle should be adapted to the pasture and to un-medicated mineral supplement before using **ULTRALYX® Free Choice Rumensin® 1620 Mineral**. Do not allow horses or other equines access to feeds containing monensin. Ingestions of monensin by horses has been fatal. Monensin medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions as reduced average daily gains may result. Do not feed to lactating goats. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: A withdrawal period has not been established in pre-ruminating calves. Do not use in calves to be processed for veal.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

Manufactured by
Ridley Block Operations
Mankato, MN 56001
www.ultralyx.com

NET WT 50 LB (22.7 kg)

Batch #

10507
UL2304
10/13