

4% Phos with IGR

Prevents adult horn fly emergence from manure of treated cattle. A nutritional supplement containing S-Methoprene* Insect Growth Regulator for continuous feeding to cattle through the horn fly season.

Active In	

S-Methoprene (CAS# 65733-16-6)	0.01%
Other Ingredients	
TOTAL	
* As the Insect Growth Regulator ALTOSID®	

Guaranteed Analysis

Guaranteed Analysis	
Calcium (Min)	18.00%
Calcium (Max)	21.60%
Phosphorus (Min)	4.00%
Salt (Min)	13.60%
Salt (Max)	16.30%
Magnesium (Min)	2.00%
Copper (Min)	1,500 ppm
Iodine (Min)	
Manganese (Min)	4,000 ppm
Selenium (Min)	26 ppm
Zinc (Min)	
Vitamin A (Min)	200,000 İU/lb
Vitamin D-3 (Min)	20,000 IU/lb
Vitamin E (Min)	20 IU/lb

Feed Ingredients

Calcium Carbonate, Monocalcium Phosphate, Dicalcium Phosphate, Salt, Molasses Products, Processed Grain By-Products, Magnesium Oxide, Hydrogenated Soybean Oil, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Sodium Selenite, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil and Red Iron Oxide.

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE REVERSE FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND FEEDING DIRECTIONS

EPA Reg No. 2724-494-7698 EPA Est. No. 7698-KY-002

NET WEIGHT: 50 LBS. (22.7 KG)

Batch #

Manufactured By: Ridlev USA Inc.

4% Phos with IGR

Prevents adult horn fly emergence from manure of treated cattle. A nutritional supplement containing S-Methoprene* Insect Growth Regulator for continuous feeding to cattle through the horn fly season.

Active Ingredient

S-Methoprene (CAS# 65733-16-6)	0.01%
Other Ingredients	
TOTAL	100.00%
* As the Insect Growth Regulator ALTOSID®	

Guaranteed Analysis

oudi di licou i li lai yolo	
Calcium (Min)	18.00%
Calcium (Max)	21.60%
Phosphorus (Min)	4.00%
Salt (Min)	13.60%
Salt (Max)	16.30%
Magnesium (Min)	
Copper (Min)	1,500 ppm
lodine (Min)	45 ppm
Manganese (Min)	4,000 ppm
Selenium (Min)	26 ppm
Zinc (Min)	
Vitamin A (Min)	
Vitamin D-3 (Min)	
Vitamin E (Min)	20 IU/lb
()	

Feed Ingredients

Calcium Carbonate, Monocalcium Phosphate, Dicalcium Phosphate, Salt, Molasses Products, Processed Grain By-Products, Magnesium Oxide, Hydrogenated Soybean Oil, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Sodium Selenite, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil and Red Iron Oxide.

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE REVERSE FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND FEEDING DIRECTIONS

EPA Reg No. 2724-494-7698 EPA Est. No. 7698-KY-002

NET WEIGHT: 50 LBS. (22.7 KG)

Batch #

Manufactured By: Ridley USA Inc.

111 West Cherry Street, Suite 500, Mankato, MN 56001

111 West Cherry Street, Suite 500, Mankato, MN 56001

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

This product contains high levels of copper. Consumption of this product by sheep or goats may result in copper toxicity. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID		
If in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical		

treatment information.

ENVIRONMENTAL HAZARDS: Do not apply directly to water, or to areas where

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FEEDING AND MANAGEMENT INSTRUCTIONS:

surface water is present, or to intertidal areas below mean high water mark.

When used as directed, **ULTRALYX® 4% Phos with IGR** will provide sufficient S-methoprene insect growth regulator to prevent the emergence of adult horn flies from manure of treated cattle. Existing adult horn flies will not be affected. Start feeding before horn flies appear and continue use until cold weather marks the end of the horn fly season. Allow free choice to cattle. Cattle should consume an average of 0.4 ounces of this product per 100 lbs of bodyweight per day, which is equivalent to 4 ounces per day for a 1000-lb. animal.

- Assure cattle are not starved for nutrients in ULTRALYX[®] 4% Phos with IGR by providing them prior to feeding ULTRALYX[®] 4% Phos with IGR.
- Offer cattle 50 pounds for every 15 head of cattle.
- 3. Remove all other self-limiting nutritional supplements.
- Locate ULTRALYX® 4% Phos with IGR where cattle congregate (watering, loafing, shade areas).
- 5. Replenish **ULTRALYX® 4% Phos with IGR** on a regular basis as needed.
- If intake of ULTRALYX® 4% Phos with IGR is below 0.4 oz. per 100 lbs. animal body weight per day, increase the number of feeding locations and/or change locations to areas more frequented by cattle.
- If intake of ULTRALYX® 4% Phos with IGR is above 0.4 oz. per 100 lbs. animal body weight per day, reduce the number of feeding locations and/or change locations to areas less frequented by cattle.
- Introduction of this product after adult horn fly infestation is established will require treatment of cattle with adulticides if elimination of the adult fly population is desired.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in a cool place. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility. **CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

EPA Reg No. 2724-494-7698 EPA Est. No. 7698-KY-002

NET WEIGHT: 50 LBS. (22.7 KG)

Altosid* is a registered trademark of Wellmark International ULTRALYX* is a registered trademark of Ridley Block Operations

Batch #

Manufactured By: **Ridley USA Inc.** 111 West Cherry Street, Suite 500, Mankato, MN 56001 34251

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING

FIRST AID

This product contains high levels of copper. Consumption of this product by sheep or goats may result in copper toxicity. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

If in eyes	1.	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	-	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
		Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below mean high water mark.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FEEDING AND MANAGEMENT INSTRUCTIONS:

When used as directed, **ULTRALYX® 4% Phos with IGR** will provide sufficient S-methoprene insect growth regulator to prevent the emergence of adult horn flies from manure of treated cattle. Existing adult horn flies will not be affected. Start feeding before horn flies appear and continue use until cold weather marks the end of the horn fly season. Allow free choice to cattle. Cattle should consume an average of 0.4 ounces of this product per 100 lbs of bodyweight per day, which is equivalent to 4 ounces per day for a 1000-lb. animal.

- Assure cattle are not starved for nutrients in ULTRALYX[®] 4% Phos with IGR by providing them prior to feeding ULTRALYX[®] 4% Phos with IGR.
- Offer cattle 50 pounds for every 15 head of cattle.
- 3. Remove all other self-limiting nutritional supplements.
- Locate ULTRALYX® 4% Phos with IGR where cattle congregate (watering, loafing, shade areas).
- 5. Replenish **ULTRALYX® 4% Phos with IGR** on a regular basis as needed.
- If intake of ULTRALYX® 4% Phos with IGR is below 0.4 oz. per 100 lbs. animal body weight per day, increase the number of feeding locations and/or change locations to areas more frequented by cattle.
- If intake of ULTRALYX® 4% Phos with IGR is above 0.4 oz. per 100 lbs. animal body weight per day, reduce the number of feeding locations and/or change locations to areas less frequented by cattle.
- Introduction of this product after adult horn fly infestation is established will require treatment of cattle with adulticides if elimination of the adult fly population is desired.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in a cool place. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility. **CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

EPA Reg No. 2724-494-7698 EPA Est. No. 7698-KY-002

NET WEIGHT: 50 LBS. (22.7 KG)

Altosid® is a registered trademark of Wellmark International ULTRALYX® is a registered trademark of Ridley Block Operations

Batch #

Manufactured By:
Ridley USA Inc.
111 West Cherry Street, Suite 500, Mankato, MN 56001

34251

Suite 500 Mankato MN 56001