



Trace Mineral Salt Block

For Beef Cattle and Horses on Pasture

GUARANTEED ANALYSIS

Calcium (Min)	8.00%
Calcium (Max)	9.60%
Phosphorus (Min)	1.00%
Salt (Min)	60.00%
Salt (Max)	65.00%
Cobalt (Min)	10 ppm
Copper (Min)	1,100 ppm
Iodine (Min)	50 ppm
Manganese (Min)	4,700 ppm
Selenium (Min)	20 ppm
Zinc (Min)	3,500 ppm
Vitamin A (Min)	20,000 IU/lb
Vitamin D-3 (Min)	2,000 IU/lb

CAUTION: Use as directed.

Consumption of selenium should not exceed 3 mg per head daily.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

INGREDIENTS

Salt, Calcium Carbonate, Molasses Products, Roughage Products, Calcium Hydroxide, Monocalcium Phosphate, Dicalcium Phosphate, Hydrogenated Soybean Oil, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Mineral Oil and Red Iron Oxide.

INTRODUCTORY PERIOD

Position the blocks 50 to 70 feet from the loafing, feeding and watering areas. Some repositioning of the blocks may be necessary as the animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption, further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of the blocks does not correct the situation, remove the blocks and evaluate the overall feeding program.

NET WEIGHT: 33.3 LB (15.12 KG)

FEEDING RECOMMENDATIONS

1. Feed at a rate of one block per 5 to 15 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing and watering areas. Adequate distance must be maintained between blocks to prevent crowding - 10 feet minimum.
3. Do not allow animals to run out of blocks. When a block is two-thirds consumed, provide a new block near it. Place very small pieces in a feed pan or trough to be cleaned up.
4. Cattle and horses should consume 2 to 4 ounces per head daily.
5. Blocks can be fed to animals in confinement. Exact location of blocks to obtain desired consumption will vary between confinement feeding programs.
6. Feed as the only self-fed sources of salt, minerals and vitamins. Provide a source of clean, fresh water at all times.

For additional information, please contact your Triple Trust® representative.



Manufactured By: **Ridley Block Operations** • Mankato, MN 56001

56100

0816