



Precalving & Rebreeding Tub with Mineralate™ Chelates

Guaranteed Analysis

Crude Protein, min.....	20.00%
Crude Fat, min.....	5.50%
Crude Fiber, max.....	3.75%
Calcium, min.....	4.50%
Calcium, max.....	5.50%
Phosphorus, min.....	1.40%
Salt, min.....	4.80%
Salt, max.....	5.80%
Magnesium, min.....	4.00%
Copper, min.....	400 ppm
Manganese, min.....	1200 ppm
Selenium, min.....	6.6 ppm
Zinc, min.....	1600 ppm
Vitamin A, min.....	50,000 IU/lb
Vitamin D3, min.....	11,000 IU/lb
Vitamin E, min.....	80 IU/lb

Ingredients

Molasses Products, Processed Grain By-Products, Monocalcium Phosphate, Dicalcium Phosphate, Salt, Calcium Carbonate, Magnesium Oxide, Calcium Hydroxide, Potassium Chloride, Zinc Amino Acid Chelate, Manganese Amino Acid Chelate, Copper Amino Acid Chelate, Magnesium Amino Acid Chelate, Potassium Amino Acid Complex, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Basic Copper Chloride, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement and Vitamin E Supplement.

Start-up feeding: Following a 7-to-10-day acclimation period, average consumption should be 0.75 to 1 pound per head per day for mature cattle with unrestricted forage supplies. Position the blocks 50 to 75 feet from loafing, feeding, and watering areas. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks, and evaluate the overall feeding program.

Feeding directions: Provide 1 block per 10 to 20 head. All animals should have equal access to blocks. Timid and smaller animals need their share. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing, and watering areas. Adequate distance must be maintained between blocks to minimize crowding - 10 feet minimum. Feed blocks continuously along with a plentiful source of forage and clean, fresh water. Cattle should not run out of blocks. When a block is two-thirds consumed, provide a new block near it. Cattle should consume an average of 0.75 to 1 pound per head daily based on a 1,000-pound animal. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding, and watering areas. If additional mineral and vitamin fortification is desired, provide an appropriate free choice mineral supplement in addition to this supplement.

CAUTION: USE AS DIRECTED. Consumption of selenium should not exceed 3 mg per head daily. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

TT3203

NET WEIGHT: 200 lbs. (90.7 kg)

09/24

**Manufactured by:
Triple Trust Tubs
Mt. Sterling, KY 40353**